

Solo-Jec[®] 5

Canine Vaccine

Description

Canine distemper-adenovirus type 2-parainfluenza-parvovirus vaccine modified live virus. Solo-Jec[®] 5 vaccine is a combination of antigenic, attenuated strains of canine distemper, parainfluenza, CAV-2 and parvovirus propagated in cell line tissue cultures. The liquid diluent is parvovirus vaccine.

Features

- Helps protect against five canine diseases
- Convenient as a single dose package with a syringe
- Contains canine parvovirus 2b and canine adenovirus type-2 (CAV-2)
- Intramuscular or subcutaneous administration

Benefits

- Convenient, ready-to-use package
- Helps protect from multiple diseases
- Safe for puppies as young as 6 weeks of age

Advantages

Helps protect against five highly contagious canine diseases in a 1 mL dose. Use in puppies as young as 6 weeks of age and adult dogs that are not exposed to *Leptospira*.

Dosage & Administration

The dosage is 1 mL injected intramuscularly or subcutaneously. Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Prepare the vaccine by injecting the diluent into the vial containing the desiccated vaccine cake. Shake well. Withdraw entire contents into the syringe. Thoroughly prepare injection site with antiseptic. Lift the loose skin behind the neck or behind the front leg and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel. Inject 1 mL intramuscularly or subcutaneously.

Administer a second dose 3 to 4 weeks later. The need for annual booster vaccination has not been established for this product. For advice on revaccination frequency, contact your veterinarian or the manufacturer.

Precautions

Store out of direct sunlight at 35°-45°F (2°-7°C). Use entire contents when first opened. Inactivate unused contents before disposal. Allergic (anaphylactoid) reactions may occur. In case of anaphylactoid reaction, administer epinephrine. Do not freeze.

Uses

Solo-Jec[®] 5 is recommended for the vaccination of healthy dogs and puppies as an aid in the reduction of disease caused by canine distemper, canine adenovirus types 1 & 2, parainfluenza and parvovirus.

Warning

Do not vaccinate pregnant animals. Do not use in ferrets or minks. Protective immunity may not be completely established in all puppies vaccinated at less than 16 weeks of age because of maternal antibody interference. Vaccinate only healthy animals. Animals incubating any disease or stressed due to shipping, malnutrition or parasitism may not achieve or maintain an adequate immune response.

Product Information

Name	Code	Size
Solo-Jec [®] 5	108842	25x1 display
Solo-Jec [®] 5	108847	25x1 tray

Highlights

- Convenient, ready-to-use package
- Helps protect puppies against five canine diseases
- Effective parvovirus, distemper and respiratory protection
- Intramuscular or subcutaneous administration



Canine Distemper-Adenovirus Type 2-Parainfluenza-Parvovirus Vaccine

Modified Live Virus

This package contains one (1 dose) vial of dry vaccine, one (1 mL) vial of vaccine diluent, and one disposable syringe.

For use in dogs only

Boehringer Ingelheim
Animal Health USA Inc.
St. Joseph, MO 64506
Phone: 1 (888) 637-4251
VLN/PCN 124/13D1.25

Indications: This product has been shown to be effective for the vaccination of healthy dogs and puppies against canine distemper virus (CDV), canine adenovirus types 1 and 2 (CAV1 & CAV2), canine parainfluenza (CPV), and canine parvovirus (CPV). This product was licensed prior to the requirement to establish a minimum age for use. The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

Precautions: Do not vaccinate pregnant animals. Under no circumstances is this product recommended for use in ferrets or mink. Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products, except as specified on this label.

See reverse side for additional instructions and precautions.

SOLO-JEC® 5
5-way


Composition: Solo-Jec® 5 vaccine is a combination of immunogenic, attenuated strains of CDV, CAV-2, CPV, and CPV type 2b, propagated in cell line tissue cultures. The CAV-2 fraction cross-protects against disease due to CAV-1. Amphotericin B, gentamicin, and thimerosal added as preservatives.

Directions and dosage: Open the syringe by twisting or tapping the cap against a hard surface to break the heat weld. Aseptically rehydrate the vaccine cake with 1 mL of vaccine diluent supplied. Shake well. Withdraw entire contents into the syringe. Administer one 1 mL dose intramuscularly or subcutaneously.

Subcutaneous administration: Lift the loose skin behind the neck or behind the front leg and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel (see note below). **Intramuscular administration:** Insert needle into muscle of the hind limb (see illustration). Inject entire contents of syringe. Do not inject directly into blood vessel. *Important note: Before injecting vaccine, pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site.* Administer a second dose 3 to 4 weeks later. The presence of maternal antibody is known to interfere with the development of active immunity in dogs and additional boosters will be required in most young animals. The need for annual booster vaccination has not been established for this product. For advice on revaccination frequency, contact your veterinarian or the manufacturer.

Precautions: Only vaccinate healthy animals. Animals incubating any disease or stressed due to shipping, malnutrition, or parasitism may not achieve or maintain an adequate immune response. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician. Inactivate unused contents before disposal.

Solo-Jec® is a registered trademark of Boehringer Ingelheim Animal Health USA Inc.



Administration sites.



123901-04

Ser:
Exp.:



Boehringer Ingelheim Animal Health USA Inc.
3239 Satellite Blvd NW | Duluth, GA 30096
Information 800-325-9167 | Sales Service 800-325-9167 | Fax 816-326-2717

SOLO-JEC® is a registered trademark of Boehringer Ingelheim Animal Health USA Inc. © 2024 Boehringer Ingelheim Animal Health USA Inc., Duluth, GA. All rights reserved. US-PET-0582-2024