# Solo-Jec<sup>®</sup> 6

# **Canine Vaccine**

# Description

Combination of immunogenic, attenuated strains of canine distemper, canine adenovirus type-2 (CAV-2), canine parainfluenza, and canine parvovirus type 2b, propagated in cell line tissue cultures. The diluent contains killed antigens for canine coronavirus vaccine. The CAV-2 fraction cross-protects against respiratory infection caused by infectious canine hepatitis (CAV-1). The CPV 2b strain has been demonstrated to be effective against CPV 2c in puppies with CPV maternal antibody.

#### Features

- Helps protect against six canine diseases in one vaccine, including coronavirus
- Convenient single dose package with syringe
- Helps protect against canine parvovirus type 2b and canine adenovirus type-2 (CAV-2)
- Helps protect against coronavirus

### Benefits

- Convenient, ready-to-use vaccine
- Helps protect against multiple diseases in one vaccine

# Advantages

Helps protect against six diseases for puppies and dogs, including coronavirus, which may cause intestinal disease in puppies.

#### Dosage & Administration

1 mL injected subcutaneously. Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Prepare the vaccine by injecting the diluent into the vial containing the desiccated vaccine cake. Shake well. Withdraw entire contents into the syringe. Thoroughly prepare injection site with antiseptic. Lift the loose skin behind the neck or behind the front leg and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel.

Administer a second dose 3 to 4 weeks later. The need for annual booster vaccination has not been established for this product. For advice on revaccination frequency, contact your veterinarian or the manufacturer.

### **Precautions**

Store out of direct sunlight at 35°-45°F (2°-7°C). Use entire contents when first opened. Inactivate unused contents before disposal. Allergic (anaphylactoid) reactions may occur. In case of anaphylactoid reaction, administer epinephrine. Do not mix with other products, except as specified on this label. Do not freeze.

# Uses

This product has been shown to be effective for the vaccination of healthy dogs 6 weeks of age or older against canine distemper virus (CDV), infectious canine hepatitis (CAV-1), canine adenovirus type-2 (CAV-2), canine coronavirus (CCV), canine parainfluenza (CPI), and canine parvovirus (CPV).

# Warning

Only vaccinate healthy animals. Animals incubating any disease or stressed due to shipping, malnutrition, or parasitism may not achieve or maintain an adequate immune response.

# **Product Information**

Name	Code	Size
Solo-Jec® 6	106085	1 mL

# Highlights

- Convenient, ready-to-use package
  Helps protect against
- 6 canine diseases in one vaccine
- Helps protect against coronavirus which may cause intestinal disease in puppies



#### **Canine Distemper-**Adenovirus Type 2-Coronavirus-Parainfluenza-**Parvovirus Vaccine** Modified Live and Killed Virus

This package contains one (1 dose) vial of dry vaccine, one (1 mL) vial of vaccine diluent, and one disposable syringe.

#### For use in dogs only

**Boehringer Ingelheim** Animal Health USA Inc. St. Joseph, MO 64506 Phone: 1 (888) 637-4251 VLN/PCN 124/1599.24

Indications: This product has been shown to be effective for the vaccination of healthy dogs 6 weeks of age or older against canine distemper virus (CDV), infectious canine hepatitis (CAV1), canine adenovirus type 2 (CAV2). canine coronavirus (CCV), canine parainfluenza (CPI), and canine parvovirus (CPV). The duration of immunity has not been determined. For more information regarding efficacy and safety data go to productdata. aphis.usda.gov.


See reverse side for instructions and precautions.

Precautions: Store out of direct sunlight at 35-46°F (2-8°C). Do not freeze. Do not mix with other products, except as specified on this label. See reverse side for additional precautions.

Preservatives: Amphotericin B. gentamicin, and thimerosal



5-way + Coronavirus



Directions and dosage: Open the syringe by twisting or tapping the cap against a hard surface to break the heat weld. Aseptically rehydrate the vaccine cake with 1 mL of vaccine diluent supplied. Shake well. Withdraw entire contents into the syringe. Thoroughly prepare the injection site with antiseptic. Administer one 1 mL dose subcutaneously: lift the loose skin behind the neck or behind the front leg and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel. Important note: Before injection site. Administer a second dose 2 to 3 weeks later. The presence of maternal antibody is known to interfere with the development of active immunity in dogs and additional boosters will be required in most young animals. The need for annual booster vaccination has not been established for this product. For advice on revaccination frequency, contact voor veterinarian or the manufacturer. your veterinarian or the manufacturer.

Precautions: Only vaccinate healthy animals. Animals incubating any disease or stressed due to shipping, malnutrition, or parasitism may not achieve or maintain an adequate immune response. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant present of a strength of the strength animals. In case of human exposure, contact a physician. Inactivate unused contents before disposal

Solo-Jec\* is a registered trademark of Boehringer Ingelheim Animal Health USA Inc.





50816-00

Ser.: Exp.:

Boehringer Ingelheim

Boehringer Ingelheim Animal Health USA Inc.

3239 Satellite Blvd NW | Duluth, GA 30096 Information 800-325-9167 | Sales Service 800-325-9167 | Fax 816-326-2717

SOLO-JEC<sup>®</sup> is a registered trademark of Boehringer Ingelheim Animal Health USA Inc. © 2024 Boehringer Ingelheim Animal Health USA Inc., Duluth, GA. All rights reserved. US-PET-0595-2024